

Table S1. Types of microorganisms isolated from amniotic fluid of women with preterm labor and preterm premature rupture of membranes.

Microorganism	Number of cases
Preterm labor and intact membranes	
<i>Ureaplasma urealyticum</i>	30
<i>Mycoplasma hominis</i>	27
Gram-positive cocci	1
<i>Lactobacillus</i> species	1
<i>Candida</i> species	1
Preterm premature rupture of membranes	
<i>Ureaplasma urealyticum</i>	37
<i>Mycoplasma hominis</i>	23
<i>Streptococci viridans</i>	2
<i>Lactobacillus</i> species	2
Gram-positive cocci	1
Unidentified gram-negative rods	1
<i>Staphylococcus aureus</i>	1
<i>Streptococcus mitis</i>	1
<i>Streptococcus agalactiae</i>	1
<i>Escherichia coli</i>	1
<i>Peptostreptococcus</i> species	1
Gram-negative rods	1

Table S2. Pairwise Spearman's rank correlation coefficients for endoglin, endostatin, IGFBP-2, IL-6, and IL-8 levels in the amniotic fluid.

	AF endoglin	AF endostatin	AF IGFBP-2	AF IL-6	AF IL-8	GA at birth
AF endostatin	0.546 ^a					
AF IGFBP-2	0.390 ^a	0.397 ^a				
AF IL-6	-0.059	0.262 ^a	-0.032			
AF IL-8	-0.152 ^c	0.222 ^b	-0.117	0.896 ^a		
GA at birth	-0.442 ^a	-0.378 ^a	-0.089	-0.344 ^a	-0.313 ^a	
GA at amniocentesis	-0.485 ^a	-0.329 ^a	-0.165 ^c	-0.114	-0.070	0.813 ^a

AF, amniotic fluid; IGFBP, insulin-like growth factor-binding protein; IL, interleukin; GA, gestational age.

^a p < 0.001

^b p < 0.01

^c p < 0.05