

Title: Combined Targeted Analysis of Metabolites and Proteins in Tear Fluid with Regard to Clinical Applications


Journal: TVST

Authors: Sascha Dammeier, Peter Martus, Franziska Klose, Michael Seid, Dario Bosch, Janina D'Alvise, Focke Ziemssen, Spyridon Dimopoulos and Marius Ueffing


Corresponding Author: Sascha Dammeier, Institute for Ophthalmic Research, Core Facility for Medical Bioanalytics, University of Tübingen, Elfriede-Aulhorn-Strasse 7, 72076 Tübingen, Germany, email: sascha.dammeier@uni-tuebingen.de

Supplementary Figure S1. Relative analyte concentrations in tear fluid of three different subjects considering the individual eyes. Data of subject 1 (mean tear secretion rates: left 1.33±0.19 mm/min, right 2.18±0.14 mm/min), subject 2 (mean tear secretion rates: left 56.67±6.60 mm/min, right 68.67±15.52 mm/min), and subject 3 (mean tear secretion rates: left 3.67±0.25 mm/min, right 3.67±0.25 mm/min) are shown for (A) amino acids, (B) acyl carnitines, (C) proteins, (D) lyso-phosphatidylcholines, (E) diacyl-phosphatidylcholines, and (F) sphingomyelins at three different visits


A



B




C




Supplementary Figure S1. continued

D



E



F

