

Table S2. Comparison of Relative Retinal Layer Thickness

Thickness (%)	Control	Post-PRP PDR	<i>P</i> value
Fovea			
NFL	6.17% ± 0.47%	7.59% ± 2.27%	0.048
GC/IPL	13.64% ± 3.50%	12.66% ± 2.88%	0.397
INL	8.01% ± 2.06%	10.49% ± 3.00%	0.02
OP/ONL	48.48% ± 3.20%	47.65% ± 3.50%	0.515
IS/OS	13.14% ± 1.40%	11.25% ± 3.12%	0.036
RPE	10.55% ± 1.80%	10.36% ± 2.23%	0.8
Parafovea			
NFL	8.12% ± 0.55%	10.01% ± 1.94%	<0.001
GC/IPL	26.99% ± 1.35%	24.86% ± 2.31%	0.002
INL	11.7% ± 1.18%	11.66% ± 1.63%	0.938
OP/ONL	34.6% ± 1.09%	36.60% ± 2.26%	0.01
IS/OS	9.78% ± 0.67%	8.81% ± 1.54%	0.032
RPE	8.81% ± 1.02%	8.06% ± 1.10%	0.069
Perifovea			
NFL	14.04% ± 1.68%	16.57% ± 2.75%	0.009
GC/IPL	21.75% ± 1.18%	21.52% ± 1.52%	0.66
INL	10.42% ± 0.96%	10.33% ± 1.86%	0.233
OP/ONL	32.81% ± 1.71%	33.82% ± 2.13%	0.082
IS/OS	11.78% ± 0.90%	9.50% ± 1.61%	<0.001
RPE	9.20% ± 1.19%	8.27% ± 1.16	0.04

Results of the relative retinal layer thickness at the follow-up visit.